



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

me that on the 14th of August last he saw at Concordia quite a number, mostly young birds, and that Dr. C. P. Blachly has in his collection a female shot some three years ago at Manhattan.—N. S. Goss, *Topeka, Kan.*

Third Addendum to List of Birds Ascertained to Occur within Ten Miles from Point des Monts, Province of Quebec, Canada; Based Chiefly upon the Notes of Napoleon A. Comeau.—(For the original list and first and second addenda see Bull. Nutt. Ornith. Club, Vol. VII, No. 4, Oct., 1882, pp. 233-242; Vol. VIII, No. 4, Oct., 1883, p. 244; and The Auk, Vol. I, No. 3, July, 1884, p. 295.)

171. *Anorthura troglodytes hyemalis*.—A pair of Winter Wrens spent the past summer (1884) at Godbout. They were first seen July 7. This species was not observed in the Gulf by either Mr. Brewster or myself, though it is common in Newfoundland.

172. *Somateria dresseri*.—Mr. William Brewster has recently called my attention to the fact that among the skins of Eider Ducks sent me by Mr. Comeau are examples of both *S. mollissima* and *S. dresseri*.

173. *Tachypetes aquila*.—A Frigate Pelican was seen and shot at by Mr. Comeau at Godbout August 13, 1884. It had previously been seen (about the end of July) by the keeper of the lightship at Manicougan, about forty miles higher up the river.

A second specimen of the Wheatear (*Saxicola ænanthe*) was taken at Godbout, September 19, 1884, and was exhibited by Mr. Comeau at the late meeting of the American Ornithologists' Union.—C. HART MERRIAM, *Locust Grove, New York.*

Albinism.—My attention was drawn to a note in the 'Oölogist' for April last, in which the writer gives his experience in albinism and asks for an explanation of these freaks of nature. In order to air my experience, and at the same time to give a probable cause, which I would like, for the sake of possible verification, other observers to look for in the future, is the object of the present note.

True albinism is of course congenital, and is a condition in which the normal pigmentary matter is deficient in the system of the individual affected; in such cases the eyes are pink, and the skin with its appendages are white or nearly so. In the case of partial albinos, however, it is difficult; their condition can probably be explained by some circumstances occurring after birth which will account for the change in the color of the skin, such for instance as the case given by the writer in the 'Oölogist,' in which the skin had been injured on the back of a Swift, and next year the patch of white feathers indicated the situation of the injury. The same thing is familiar in the case of the horse whose back or shoulder is galled by the harness; white patches appear, owing to lowered vitality of the injured part. These cases are familiar, but I wish to give possibly another cause acting in the same way, only more general. It is this. When a boy I shot among others a black squirrel peculiarly marked, it having a per-